arXiv:math/0609621v2 [math.NT] 15 Oct 2006

On the solutions to a power sum problem

Johan Andersson[*]

February 2, 2008

Abstract

In a recent paper [2] we proved that

$$\inf_{|z_k|=1} \max_{\nu=1,\ldots,n^2-n} \left| \sum_{k=1}^{n} z_k^{\nu} \right| = \sqrt{n-1}$$

when $n-1$ is a prime power. We also proved that a construction of Fabrykowski of an explicit n-tuple $(z_1,\ldots,z_n)$ provided a global minimum to the problem. The construction depends on the construction of perfect difference sets of order $n-1$. As an open problem we asked whether we could obtain all global minima by this construction. In this paper we will show that this is in fact true and we furthermore prove that

$$\inf_{|z_k|=1} \max_{\nu=1,\ldots,n^2-n} \left| \sum_{k=1}^{n} z_k^{\nu} \right| > \sqrt{n-1}$$

in case there exist no perfect difference set of order $n-1$. A well known conjecture states that each perfect difference set must have prime power order.

1 Introduction

Throughout this paper we will let

$$S(\nu) = \sum_{k=1}^{n} z_k^{\nu} \tag{1}$$

denote the pure power sums. In the papers [1],[2] and [3] we considered Turán's problem 10 which is to estimate

$$(*) = \inf_{|z_k|\geq 1} \max_{\nu=1,\ldots,n^2} |S(\nu)|.$$

---

[*] Department of Mathematics, Stockholm University, SE-10691, Sweden. *johana@math.su.se*

We showed already in [1] that $\sqrt{n} \leq (*) \leq \sqrt{n+1}$ if $n+1$ is a prime. In general we have that $\sqrt{n} \leq (*)$ (see [1]) and $(*) = \sqrt{n} + O(n^{0.2625+\epsilon})$ (see [3]). It would be interesting to find an explicit solution (in terms of minimal systems $(z_1, \ldots, z_n)$) to this inf max problem, but due to lack of success in this we considered (see [2]) the modified problems

$$(**) = \inf_{|z_k| \geq 1} \max_{\nu=1,\ldots,n^2-2} |S(\nu)|,$$

and

$$(***) = \inf_{|z_k| = 1} \max_{\nu=1,\ldots,n^2-n} |S(\nu)|.$$

We proved that $(**) = \sqrt{n}$ if n is a prime power and $(***) = \sqrt{n-1}$ if $n-1$ is a prime power. We gave explicit solutions and asked whether these were the only solutions possible. In this paper we will answer this question in the case $(***)$ ([2], Problem 2). In our paper we used a theorem independently proved by Newman, Cassels and Szalay (Szalay [8] and Theorem 7.3 of Turán [9]) to get a lower bound, and our solution will come from examining the steps of the proof of that theorem (compare the proof of Lemma 1 below and page 81 in Turán [9]) and modifying the proof to better fit our purpose.

2 A lower bound

Lemma 1. *Suppose that $(z_1, \ldots, z_n)$ is an n-tuple of unimodular complex numbers. One then has that*

(i) $\max_{\nu=1,\ldots,n^2-n} |S(\nu)| \geq \sqrt{n-1}$.

(ii) If the inequality in (i) holds with equality, then $|S(\nu)| = \sqrt{n-1}$ for all $\nu = 1, \ldots, n^2 - n$.

Proof. Let θ_k be defined so that $z_k = e(\theta_k)$. Then

$$S(\nu) = \sum_{k=1}^{n} e(\nu\theta_k). \qquad (e(x) = e^{2\pi i x})$$

Let

$$F_m(t) = \sum_{\nu=1-m}^{m-1} \left(1 - \frac{|\nu|}{m}\right) e(\nu t)$$

be the m'th Fejér kernel. The Fejér kernel can be written as

$$F_m(t) = \frac{1}{m} \left(\frac{\sin \pi m x}{\sin \pi x}\right)^2, \tag{2}$$

and is thus non negative. We have that

$$\sum_{\nu=1-m}^{m-1}\left(1-\frac{|\nu|}{m}\right)|S(\nu)|^2=\sum_{k,l=1}^{n}F_m(\theta_k-\theta_l),$$

which by the contribution of the diagonal $k=l$, and the non negativity of the Fejér kernel implies that

$$\sum_{\nu=1-m}^{m-1}\left(1-\frac{|\nu|}{m}\right)|S(\nu)|^2\geq nF_m(0).$$

By letting $m=n^2-n+1$ and $|S(\nu)|^2=n-1+\varepsilon_\nu$ we get

$$\sum_{\nu=n-n^2}^{n^2-n}\left(1-\frac{|\nu|}{n^2-n+1}\right)\varepsilon_\nu\geq F_{n^2-n+1}(0).$$

By the fact that $|S(\nu)|=|S(\nu)|$ we have that $\varepsilon_{-\nu}=\varepsilon_\nu$. By furthermore using the facts that $F_{n^2-n+1}(0)=n^2-n+1$ and $\varepsilon_0=n^2-n+1$, we obtain the following inequality

$$\sum_{\nu=1}^{n^2-n}\left(1-\frac{\nu}{n^2-n+1}\right)\varepsilon_\nu\geq 0.$$

This implies that for at least one $1\leq\nu\leq n^2-n$ we have that $\varepsilon_\nu\geq 0$, which implies (i). Furthermore if all $\varepsilon_\nu\leq 0$ then $\varepsilon_\nu=0$ for all $\nu=1,\ldots,k$ which implies (ii) ☐

Lemma 2. *Let $(z_1,\ldots,z_n)$ be an n-tuple of unimodular complex numbers. Then the following statements are equivalent:*

(i) $S(\nu)=0$ for $\nu=1,\ldots,n-1$.

(ii) The numbers z_k are the n vertices of a regular n-gon on the unit circle.

Proof. The fact that $(ii)\implies(i)$ is immediate. Remains to prove that $(i)\implies(ii)$. Consider the polynomial

$$p(x)=\prod_{k=1}^{n}(x-z_k)=x^n+a_1x^{n-1}+\cdots+a_n.$$

By the Newton-Girard identities

$$S(\nu)+a_1S(\nu-1)+\cdots+a_{\nu-1}S(1)+\nu a_\nu=0,\qquad(\nu=1,2,\ldots,n)$$

and the fact that $S(\nu)=0$ for $\nu=1,\ldots,n-1$ we find that $a_\nu=0$ for $\nu=1,\ldots,n-1$. We see that

$$p(x)=x^n+(-z_1)\cdots(-z_n)=x^n-w$$

for some complex number w with $|w|=1$. ☐

3 Perfect Difference sets

Lemma 3. (Singer [7]) *Let q be a prime power. Then there exists a perfect difference set of order q or in other words integers $a_1, \ldots, a_{q+1}$ such that the integers $a_i - a_j$ for $i \neq j$ form all non zero residues* $\mod q^2 + q + 1$.

Proof. This follows by the use of projective planes over finite fields (See Singer [7]). □

Conjecture 1. (The prime power conjecture) *There only exist perfect different sets of order q if q is a prime power.*

The prime power conjecture has been proved for a lot of special cases. If $n \equiv 1, 2 \pmod 4$ and can not be written as a sum of two squares then there exist no perfect difference set of order n (Bruck-Ryser [5]). In case $n \geq 6$ and $n \equiv 3, 6 \pmod 9$ (Wilbrink [10]). Finally it has been proved for $n < 2 \cdot 10^9$ (Baumert-Gordon [4]). We remark here that the problem that a general finite projective plane has prime power order is more difficult (The Bruck-Ryser result is still true, but other methods fails), and is yet to be proved true for $n = 12$. Conjecture 1 is equivalent to the conjecture that a finite *cyclic* projective plane must have prime power order.

4 Main result

Theorem 1. *Let $(z_1, \ldots, z_n)$ be an n-tuple of unimodular complex numbers. Then the following statements are equivalent:*

(i) $|S(\nu)| = \sqrt{n-1}$ *for* $\nu = 1, \ldots, n^2 - n$.

(ii) There exist a complex number α with $|\alpha| = 1$ and a perfect difference set $(a_1, \ldots, a_n)$ of order $n - 1$ (which means that $a_i - a_j$ for $i \neq j$ form all non zero residues $\mod n^2 - n + 1$*) such that* $z_k = \alpha\, e\left(\dfrac{a_k}{n^2 - n + 1}\right)$.

In case (i) and (ii) are false then one has that $\max\limits_{\nu=1,\ldots,n^2-n} |S(\nu)| > \sqrt{n-1}$.

Proof. By choosing

$$z_k = \alpha e(\theta_k), \qquad \big(e(x) = e^{2\pi i x}\big)$$

with

$$\theta_k = \frac{a_k}{n^2 - n + 1}, \qquad \theta_1 = 0 \qquad \text{and} \qquad |\alpha| = 1,$$

we see that

$$\begin{aligned}\left|\sum_{k=1}^{n} z_k^{\nu}\right|^2 &= n + \sum_{k\neq l} e(\nu(\theta_k - \theta_l)),\\ &= n - 1 + \sum_{j=1}^{n^2-n+1} e(\nu\lambda_j),\end{aligned}$$

where $\lambda_1 = 0$ and $0 \leq \lambda_2 \leq \cdots \leq \lambda_{n^2-n+1} \leq 1$ consists of the $n^2 - n$ differences $\theta_k - \theta_l$ where $k \neq l$. Then

$$|S(\nu)| = \sqrt{n-1} \qquad\qquad (\nu = 1, \ldots, n^2 - n)$$

is equivalent to

$$\sum_{j=1}^{n^2-n+1} e(\nu\lambda_j) = 0. \qquad\qquad (\nu = 1, \ldots, n^2 - n)$$

By Lemma 2 this is true iff λ_j consists of the vertices of a regular $(n^2 - n + 1)$-gon. Since $\lambda_0 = 1$ and λ_j is increasing this is true iff $\lambda_j = \dfrac{j}{n^2-n+1}$. This is equivalent to the fact that $a_i - a_j$ for $i \neq j$ form all non zero residues mod $n^2 - n + 1$. That the a_j's are furthermore integers follows by the fact that $\theta_k = \theta_k - 0 = \theta_k - \theta_0 = \dfrac{j_k}{n^2-n+1}$ for integers j_k. This means that $(a_1, \ldots, a_n)$ form a perfect difference set of order $n - 1$. In case $|S(\nu)| \neq \sqrt{n-1}$ for some $1 \leq \nu \leq n^2 - n$ then it follows from Lemma 1 that $|S(\nu)| > \sqrt{n-1}$ for some $1 \leq \nu \leq n^2 - n$. □

As consequences of our theorem we have the following corollaries:

Corollary 1. *Suppose that there exist no perfect difference set of order* $n - 1$*. Then one has that*

$$\inf_{|z_k|=1} \max_{\nu=1,\ldots,n^2-n} |S(\nu)| > \sqrt{n-1}.$$

Corollary 2. *The prime power conjecture is equivalent to the following statement: Suppose that* $n - 1$ *is not a prime power. Then*

$$\inf_{|z_k|=1} \max_{\nu=1,\ldots,n^2-n} |S(\nu)| > \sqrt{n-1}.$$

Corollary 3. *Suppose that $n-1$ is a prime power. Then one has that*

$$\inf_{|z_k|=1} \max_{\nu=1,\ldots,n^2-n} |S(\nu)| = \sqrt{n-1},$$

and every minimal system is given by the construction $z_k = \alpha e\left(\frac{a_k}{n^2-n+1}\right)$ *of Fabrykowski with* $|\alpha| = 1$ *and where* $(a_1, \ldots, a_n)$ *is a perfect difference set of order* $n-1$.

Proof. This follows from Theorem 1 and Lemma 3 with $q = n-1$. Further discussion of the Fabrykowski construction is given in [6] and [2]. ☐

References

[1] J. Andersson. On some power sum problems of Turán and Erdős. *Acta Math. Hungar.*, 70(4):305–316, 1996.

[2] ——— Explicit solutions to certain inf max problems from Turan power sum theory, 2006. arXiv:math.NT/0607238.

[3] ——— Turán's problem 10 revisited. arXiv:math.NT/0609271.

[4] Leonard D. Baumert and Daniel M. Gordon. On the existence of cyclic difference sets with small parameters. In *High primes and misdemeanours: lectures in honour of the 60th birthday of Hugh Cowie Williams*, volume 41 of *Fields Inst. Commun.*, pages 61–68. Amer. Math. Soc., Providence, RI, 2004.

[5] R. H. Bruck and H. J. Ryser. The nonexistence of certain finite projective planes. *Canadian J. Math.*, 1:88–93, 1949.

[6] J. Fabrykowski. A note on sums of powers of complex numbers. *Acta Math. Hungar.*, 62(3-4):209–210, 1993.

[7] J. Singer. A theorem in finite projective geometry and some applications to number theory. *Trans. Amer. Math. Soc.*, 43(3):377–385, 1938.

[8] M. Szalay. *On number theoretical extremal problems.* Thesis, 1974.

[9] P. Turán *On a new method of analysis and its applications.* Pure and Applied Mathematics (New York). John Wiley & Sons Inc., New York, 1984. With the assistance of G. Halász and J. Pintz, With a foreword by Vera T. Sós, A Wiley-Interscience Publication.

[10] H. A. Wilbrink. A note on planar difference sets. *J. Combin. Theory Ser. A*, 38(1):94–95, 1985.